

228-313

6/18/2014

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Danielle A. Larochele  
Nufarm Americas, Inc.  
4020 Aerial Center Parkway, Suite 101  
Morrisville, NC 27560

JUN 18 2014

Subject: Notification; Per PR-Notice 98-10  
HORSEPOWER SELECTIVE HERBICIDE  
EPA Reg. No. 228-313  
Date Submitted: June 13, 2014

Dear Ms. Larochele:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated June 13, 2014 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact Kable Bo Davis at (703) 306-0415 or [davis.kable@epa.gov](mailto:davis.kable@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "K. Montague".

Kathryn Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

2/9

# NOTIFICATION

Please read instructions on reverse before completing form. Approved, OMB No. 2070-0060, Approval expires 05-31-98

|   |  |                          |
|---|--|--------------------------|
| <b>EPA</b><br>United States<br><b>Environmental Protection Agency</b><br>Washington, DC 20460 | <input type="checkbox"/> <b>Registration</b><br><input checked="" type="checkbox"/> <b>Amendment</b><br><input type="checkbox"/> <b>Other:</b> | OPP Identifier<br>Number |
|---|--|--------------------------|

### Application for Pesticide - Section I

|   |  |  |
|---|--|--|
| <b>1. Company/Product Number</b><br>228-313   | <b>2. EPA Product Manager</b><br>Ms. Kathryn Montague  | <b>3. Proposed Classification</b><br><br><input checked="" type="checkbox"/> None<br><input type="checkbox"/> Restricted |
| <b>4. Company/Product (Name)</b><br>HORSEPOWER® Selective Herbicide   | <b>PM#</b><br>23   |  |
| <b>5. Name and Address of Applicant (Include ZIP Code)</b><br>Nufarm Americas Inc.<br>4020 Aerial Center Parkway, Suite 101<br>Morrisville, NC 27560<br><br><input type="checkbox"/> Check if this is a new address | <b>6. Expedited Review.</b><br><br>In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to:<br><br>EPA Reg. No.: _____<br>Product Name: _____ |  |

### Section - II

|  |  |
|--|--|
| <input type="checkbox"/> Amendment - Explain below.                            | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application  |
| <input checked="" type="checkbox"/> Notification - Explain below.              | <input type="checkbox"/> Other - Explain below   |

**Explanation: Use additional page(s) if necessary. (For Section I and Section II.)** **NOTIFICATION**

**NOTIFICATION OF LABEL CHANGES Per PR Notice 98-10** **JUN 18 2014**

(See attachment for certification)

**CONTACT:** Danielle Larochelle, Nufarm Americas Inc, 4020 Aerial Center Parkway, Suite 101, Morrisville, NC 27560.  
 Tel: 919-379-2530. Email: danielle.larochelle@us.nufarm.com

### Section - III

|  |   |   |   |   |  |
|--|---|---|---|---|--|
| <b>1. Material This Product Will Be Packaged In:</b>   |   |   |   | <b>2. Type of Container</b>   |  |
| <b>Child-Resistant Packaging</b><br><input type="checkbox"/> Yes*<br><input checked="" type="checkbox"/> No  | <b>Unit Packaging</b><br><input type="checkbox"/> Yes<br><input checked="" type="checkbox"/> No<br><br>If "Yes"<br>Unit Packaging wgt.      No. per container | <b>Water Soluble Packaging</b><br><input type="checkbox"/> Yes<br><input checked="" type="checkbox"/> No<br><br>If "Yes"<br>Package wgt.      No. per container | <input type="checkbox"/> Metal<br><input checked="" type="checkbox"/> Plastic<br><input type="checkbox"/> Glass<br><input type="checkbox"/> Paper<br><input type="checkbox"/> Other (Specify) _____ |   |  |
| <b>*Certification must be submitted</b>  |   |   |   |   |  |
| <b>3. Location of Net Contents Information</b><br><input checked="" type="checkbox"/> Label <input type="checkbox"/> Container   |   | <b>4. Size(s) Retail Container</b><br>1, 2.5, 30, 220, 250 Gallons  |   | <b>5. Location of Label Directions</b><br><input checked="" type="checkbox"/> On Label<br><input type="checkbox"/> On labeling accompanying product |  |
| <b>6. Manner in Which Label is Affixed to Product</b><br><input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input checked="" type="checkbox"/> Other: <u>crack and peel</u><br><input type="checkbox"/> Stenciled |   |   |   |   |  |

### Section - IV

|  |                                       |  |
|--|---------------------------------------|--|
| <b>1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)</b>  |                                       |  |
| <b>Name</b><br>Danielle A. Larochelle  | <b>Title</b><br>Regulatory Manager    | <b>Telephone No. (Include Area Code)</b><br>919-379-2530 |
| <b>Certification</b>   |                                       | <b>6. Date Application Received (Stamped)</b>            |
| I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. |                                       |  |
| <b>2. Signature</b><br>  | <b>3. Title</b><br>Regulatory Manager |  |
| <b>4. Typed Name</b><br>Danielle A. Larochelle   | <b>4. Date</b><br>June 13, 2014       |  |



4/9

# HORSEPOWER®

## SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES. CONTROLS BLACK MEDIC, CHICKWEED, CLOVER COCKLEBUR, DANDELION, KNOTWEED, OXALIS, PLANTAIN, THISTLE AND MANY OTHER SPECIES OF BROADLEAF WEEDS; SOME OF WHICH ARE LISTED ON THIS LABEL.

CONTAINS MCPA, TRICLOPYR AND DICAMBA

**ACTIVE INGREDIENTS:**

|   |        |
|---|--------|
| Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*        | 48.99% |
| Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid** | 5.59%  |
| Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***     | 4.82%  |

|                           |                |
|---------------------------|----------------|
| <b>OTHER INGREDIENTS:</b> | 40.60%         |
| <b>TOTAL</b>              | <b>100.00%</b> |

By Isomer Specific AOAC Method, Equivalent to:

|   |       |                |
|---|-------|----------------|
| *2-Methyl-4-Chlorophenoxyacetic Acid        | 40.0% | 3.80 lbs./gal. |
| **3,5,6-Trichloro-2-Pyridinyloxyacetic Acid | 4.0%  | 0.38 lbs./gal. |
| ***3,6-Dichloro-o-Anisic Acid               | 4.0%  | 0.38 lbs./gal. |

### KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

NOTE: Spanish language is optional

EPA REG. NO. 228-313  
EPA EST. NO. 228-IL-1

MANUFACTURED BY  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE 11901 S. AUSTIN AVE.  
BURR RIDGE, IL 60527 ALSIP, IL 60803



NOTIFICATION

JUN 18 2014

[Nufarm Grow a better tomorrow]  
[Grow a better tomorrow]

NET CONTENTS GALS.

5/9

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid contact with eyes, skin or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are Butyl rubber ≥1 4 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Mixers, loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants,
- Protective eyewear (goggles or face shield)\*
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof materials.

- Eye protection is not required ONLY when 3 pints of product is diluted with 20 or more gallons of water

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow people or pets to enter the treated field until sprays have dried.

**FIRST AID**

|                               |  |
|-------------------------------|--|
| <b>IF IN EYES</b>             | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |
| <b>IF SWALLOWED</b>           | <ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>IF ON SKIN OR CLOTHING</b> | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

This product is corrosive to the eyes. Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

6/9

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees\* and Roughs) . Not for use on turf being grown for resale or other commercial use as sod, or for commercial seed production or for research purposes. \*Excluding Bentgrass Tees.

**Entry Restrictions:** Do not allow people or pets to enter treated area until sprays have dried.

**Application Restrictions:** Avoid contact with eyes, skin, or clothing during and after application. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. This product may cause skin sensitization reactions in some people.

The maximum application rate is 3.2 pints of product (1.5 lb ae) per acre per year.

Limit two applications per year with a minimum retreatment interval of 21 days.

Aerial application is prohibited.

### USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperature over 90 degrees, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

**For optimum results:** (1) avoid applying during excessively dry or hot periods unless irrigation is used;(2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.


### SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Do not apply with a nozzle height greater than 4 feet above the crop canopy for ground boom application.



## WEEDS CONTROLLED

|                   |                     |                                 |  |
|-------------------|---------------------|---------------------------------|--|
| Alfalfa           | Dog fennel          | Mayweed                         | Spurweed   |
| Amaranth          | Dragonhead mint     | Milkweed                        | Starbur  |
| Aster             | Evening primrose    | Morningglory                    | Stinging nettle                                  |
| Beadstraw         | Field pennycress    | Mustard                         | Stitchwort                                       |
| Bedstraw          | Fleabane            | Nightflowering catchfly         | Stunkweed  |
| Beggartick        | Florida pusley      | Nightshade                      | Sumpweed   |
| Beggarweed        | Frenchweed          | Oxalis(Stricta and corniculata) | Sunflower  |
| Bindweed          | Galinsoga           | Parsley-piert                   | Tansy ragwort                                    |
| Bitter sneezeweed | Geranium            | Pennywort                       | Teasel   |
| Black medic       | Goatsbeard          | Peppergrass                     | Teaweed  |
| Bladder campton   | Goldenrod           | Pepperweed                      | Texas blueweed                                   |
| Bracken fern      | Goosefoot           | Pigweed                         | Thistle  |
| Broomweed         | Gromwell            | Plantain                        | Toadflax   |
| Buckhorn          | Ground ivy          | Poison hemlock                  | Trumpet creeper                                  |
| Buckwheat         | Halfshrub sundrop   | Poison ivy                      | Velvetleaf                                       |
| Buffalobur        | Hawkweed            | Poison oak                      | Veronica   |
| Burclover         | Healall             | Pokeweed                        | Vetch  |
| Burcucumber       | Heartleaf drymary   | Poorjoe                         | Waterhemlock                                     |
| Burdock           | Hemp sesbania       | Prickly sida                    | Waterhemp  |
| Bursage           | Hempnettle          | Prickly lettuce                 | Whitebrush                                       |
| Buttercup         | Henbit              | Puncturevine                    | Wild violet                                      |
| Canada thistle    | Horsenettle         | Purslane                        | Wild radish                                      |
| Carpetweed        | Horseweed           | Ragweed                         | Wild onion                                       |
| Catnip            | Ironweed            | Red sorrel                      | Wild aster                                       |
| Catsear           | Jerusalem artichoke | Red clover                      | Wild carrot                                      |
| Chamise           | Jimsonweed          | Redvine                         | Wild lettuce                                     |
| Chamomile         | Knapweed            | Redstem                         | Wild garlic                                      |
| Chickweed         | Knapwood            | Rubberweed                      | Wild geranium                                    |
| Chicory           | Knawel              | Sheep sorrel                    | Wood sorrel                                      |
| Cinquefoil        | Knotweed            | Shepherdspurse                  | Wormwood   |
| Clover            | Kochia              | Sicklepod                       | Yankeeweed                                       |
| Cocklebur         | Ladysthumb          | Smartweed                       | Yarrow   |
| Coffeeweed        | Lambsquarter        | Snakeweed                       | Yellow starthistle                               |
| Cornflower        | Lespedeza           | Sowthistle                      | Yellow rocket and many other<br>broadleaf weeds. |
| Cornspeedwell     | Little starwort     | Speedwell                       |  |
| Croton            | Mallow              | Spiderwort                      |  |
| Daisy             | Marc's Tail         | Spikeweed                       |  |
| Dandelion         | Marshelder          | Spurge                          |  |
| Dock              | Matchweed           | Spurry                          |  |

## ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

**Herbi™ Controlled Droplet Applicator:** Add 1-1/2 to 2-1/4 pints of this product to the 5 pint Herbi bottle, then fill with water to make 5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution.

While walking at approximately 1 pace (3 feet) per second, spray entire contents over 33,000 square feet (3/4 of an acre). Do not overlap (double coverage) at edge of spray patterns. Reduced rates (use 1/2 of rate shown above) of this product must be applied when grass is stressed from heat, drought, etc.

**Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators:** Use this product at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

**NOTE:** For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

8/9

### COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 1 68D, Amoco Oil's Amoco Spray Mate and Universal Co-op.'s Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PRODUCT DISPOSAL:** Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING: Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Nonrefillable containers larger than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**OR**

**Refillable containers larger than 5 gallons:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

